arXiv:1201.4985v1 [math-ph] 24 Jan 2012

Local Generalized Pauli's Theorem

Nikoly Marchuk and Dmitry Shirokov

January 25, 2012

Abstract

In this work a generalized Pauli's theorem (proved by D. S. Shirokov for two sets e^a, h^a, $a = 1, \ldots, n$ of Clifford algebra elements $\mathcal{C}\!\ell(p, q)$, $p + q = n$) is extended to the case, when one or both sets of elements depends smoothly on points x of the Euclidian space. We prove that in this case for any point x_0 of Euclidian space there exists ε-neighborhood $O_\varepsilon(x_0)$ and there exists smooth function $T : O_\varepsilon(x_0) \to \mathcal{C}\!\ell(p, q)$ such that two sets of Clifford algebra elements are connected by a similarity transformation $h^a(x) = T(x)^{-1} e^a(x) T(x)$, $a = 1, \ldots, n$ in $O_\varepsilon(x_0)$.

Let V be r-dimensional Euclidian space with scalar product (x, y), $\forall x, y \in V$ and with the norm

$$\|x\| = \sqrt{(x, x)}, \quad \forall x \in V.$$

Let Ω be domain in V and let $\varepsilon > 0$ be positive real number. ε-neighborhood of a point $x_0 \in V$ is the domain

$$O_\varepsilon(x_0) = \{x \in V \ : \ \|x - x_0\| < \varepsilon\}.$$

Consider a real or complex ($\mathbb{F} = \mathbb{R}$ or $\mathbb{F} = \mathbb{C}$) Clifford algebra $\mathcal{C}\!\ell^{\mathbb{F}}(p, q)$ of dimension[1] $n = p + q$ with generators e^a, $a = 1, \ldots, n$ and with basis of 2^n elements

$$e, e^a, e^{ab}, \ldots, e^{1\ldots n}, \tag{1}$$

---

[1] An important special case is $r = n$, when the basis of V is the set of generators e^a. In this case V can be considerd as a pseudo-Euclidian space with two metrics - Euclidian and pseudo-Euclidian. We use Euclidian metric to determine a neighborhood of a point.

enumerated by ordered multi-indices of length from 0 to n.

The generators satisfy the relations

$$e^a e^b + e^b e^a = 2\eta^{ab} e,$$

where η^{ab} are elements of the diagonal matrix η of order n with p elements equals 1 and q elements equals -1 on the diagonal.

Consider a function

$$f : \Omega \to \mathcal{C}\!\ell^{\mathbb{F}}(p,q)$$

with values in Clifford algebra $\mathcal{C}\!\ell^{\mathbb{F}}(p,q)$. The function $f = f(x)$ can be written in the form

$$f = ue + u_a e^a + \ldots + u_{1\ldots n} e^{1\ldots n},$$

where $u = u(x)$, $u_a = u_a(x)$, $\ldots$ are functions $\Omega \to \mathbb{F}$ and basis elements (1) do not depend on $x \in V$.

If functions $u, u_a, \ldots, u_{1\ldots n}$ have continuous derivatives up to order k in Ω, then we say that (real or complex) functions $u, u_a, \ldots, u_{1\ldots n}$ and f belong to the class $C^k(\Omega)$ ($C^0(\Omega)$ – the class of continuous functions in domain Ω).

Theorem 1 (Local generalized Pauli's theorem in the case of even n).

Let n be even positive number and $h^a = h^a(x)$, $a = 1, \ldots, n$ are functions $\Omega \to \mathcal{C}\!\ell^{\mathbb{F}}(p,q)$ of class $C^k(\Omega)$ such that

$$h^a(x)h^b(x) + h^b(x)h^a(x) = 2\eta^{ab} e, \quad a, b = 1, \ldots, n, \quad \forall x \in \Omega.$$

Then for any $x_0 \in \Omega$ there exists $\varepsilon > 0$ and there exists a function $T = T(x) : O_\varepsilon(x_0) \to \mathcal{C}\!\ell^{\mathbb{F}}(p,q)$, satisfying the conditions

1. *$T(x)$ – function of class $C^k(O_\varepsilon(x_0))$;*
2. *$T(x)$ – an invertible element of Clifford algebra $\mathcal{C}\!\ell^{\mathbb{F}}(p,q)$ for any $x \in O_\varepsilon(x_0)$;*
3. *$e^a = T^{-1}(x)h^a(x)T(x)$, $a = 1, \ldots, n$, $\forall x \in O_\varepsilon(x_0)$;*
4. *The function $T(x)$ is defined up to multiplication by (real in the case $\mathbb{F} = \mathbb{R}$ or complex in the case $\mathbb{F} = \mathbb{C}$) function of class $C^k(O_\varepsilon(x_0))$ that is not equal to zero for any point of $O_\varepsilon(x_0)$.*

Proof. We denote elements of the basis (1) by e^A, where A are ordered multi-indices of length from 0 to n. Let denote $e_A = (e^A)^{-1}$. For each basis element e^A we have $(e^A)^2 = e$ or $(e^A)^2 = -e$, so for all A we have $e_A = \pm e^A$.

We denote

$$\begin{aligned} h^{ab} &= h^a h^b, \quad \text{for} \quad 1 \leq a < b \leq n; \\ h^{abc} &= h^a h^b h^c, \quad \text{for} \quad 1 \leq a < b < c \leq n \end{aligned}$$

and so on. Let denote elements

$$e, h^a, h^{ab}, \ldots, h^{1\ldots n}$$

by h^A.

Let consider sum of the following form (sum over all ordered multi-indices A of length from 0 to n)

$$\sum_A h^A(x) F e_A \tag{2}$$

where F is an element of the basis (1).

Consider an arbitrary point $x_0 \in \Omega$. By generalized Pauli's theorem [1] we have at least one basis element (1) (denote it by F) for which

$$T = \sum_A h^A(x_0) F e_A \neq 0. \tag{3}$$

We define the norm of Clifford algebra elements by

$$|U| = \sqrt{\mathrm{Tr}(U^\dagger U)},$$

where the operation of Hermitian conjugation $\dagger$ is defined in [2]. Then from (3) we obtain

$$|T| = |\sum_A h^A(x_0) F e_A| = \delta > 0.$$

Since a linear combination of functions of class $C^k(\Omega)$ is a function of class $C^k(\Omega)$, then the expression $|\sum_A h^A(x) F e_A|$ is a continuous function of $x \in \Omega$. Then there exists a real number $\varepsilon > 0$ such that

$$|\sum_A h^A(x) F e_A| > \delta/2, \quad \forall x \in O_\varepsilon(x_0)$$

in ε-neighborhood of the point x_0.

Consequently, we construct a function

$$T(x) = \sum_A h^A(x) F e_A \neq 0, \quad \forall x \in O_\varepsilon(x_0)$$

of class $C^k(O_\varepsilon(x_0))$. By generalized Pauli's theorem [1] we have

$$e^a = T^{-1}(x) h^a(x) T(x), \quad a = 1, \ldots, n, \quad \forall x \in O_\varepsilon(x_0))$$

or, equivalently,

$$h^a(x) = T(x) e^a T^{-1}(x), \quad a = 1, \ldots, n, \quad \forall x \in O_\varepsilon(x_0)).$$

The theorem is proved.

The theorem evidently generalizes to the case when both sets of Clifford algebra elements depend on points x of the Euclidian space.

Let formulate and prove a similar theorem for the case of odd n.

Theorem 2 (Local generalized Pauli's theorem in the case of odd n for real Clifford algebra).

Let n be positive odd number and $h^a = h^a(x)$, $a = 1, \ldots, n$ are functions $\Omega \to \mathcal{C}\!\ell^{\mathbb{R}}(p,q)$ of class $C^k(\Omega)$ such that

$$h^a(x) h^b(x) + h^b(x) h^a(x) = 2\eta^{ab} e, \quad a, b = 1, \ldots, n, \quad \forall x \in \Omega.$$

Then the product $h^1(x) h^2(x) \ldots h^n(x) = h^{1\ldots n}$ does not depend on x and equals $\pm e^{1\ldots n}$ or $\pm e$ (last case is possible only if $p - q = 1 \mod 4$).

Then for any point $x_0 \in \Omega$ there exists $\varepsilon > 0$ and there exists a function $T = T(x) : O_\varepsilon(x_0) \to \mathcal{C}\!\ell^{\mathbb{R}}(p,q)$, satisfying the conditions

1. *$T(x)$ – function of class $C^k(O_\varepsilon(x_0))$;*
2. *$T(x)$ – an invertible element of Clifford algebra $\mathcal{C}\!\ell^{\mathbb{R}}(p,q)$ for any $x \in O_\varepsilon(x_0)$;*
3. *(a) $e^a = T^{-1}(x) h^a(x) T(x) \quad \Leftrightarrow \quad h^{1\ldots n} = e^{1\ldots n}$,*
   (b) $e^a = -T^{-1}(x) h^a(x) T(x) \quad \Leftrightarrow \quad h^{1\ldots n} = -e^{1\ldots n}$,
   (c) $e^a = e^{1\ldots n} T^{-1}(x) h^a(x) T(x) \quad \Leftrightarrow \quad h^{1\ldots n} = e$,

(d) $e^a = -e^{1\dots n}T^{-1}(x)h^a(x)T(x) \quad \Leftrightarrow \quad h^{1\dots n} = -e$,

where equalities hold for $a = 1,\dots,n$ *and* $\forall x \in O_\varepsilon(x_0)$*;*

4. *The function* $T(x)$ *is defined up to multiplication by elements* $\lambda(x)e + \nu(x)e^{1\dots n}$*, where* $\lambda(x)$ *and* $\nu(x)$ *are real functions of class* $C^k(O_\varepsilon(x_0))$ *such that element* $\lambda(x)e+\nu(x)e^{1\dots n}$ *is invertible for any point of domain* $O_\varepsilon(x_0)$*.*

Before discussing the proof of this theorem let us formulate another theorem.

Theorem 3 (Local generalized Pauli's theorem in the case of odd n for complex Clifford algebra).

Let n *be positive odd number and* $h^a = h^a(x)$*,* $a = 1,\dots,n$ *are functions* $\Omega \to \mathcal{C}\ell^{\mathbb{C}}(p,q)$ *of class* $C^k(\Omega)$ *such that*

$$h^a(x)h^b(x) + h^b(x)h^a(x) = 2\eta^{ab}e, \quad a,b = 1,\dots,n, \quad \forall x \in \Omega.$$

Then the product $h^1(x)h^2(x)\dots h^n(x) = h^{1\dots n}$ *does not depend on* x *and equals* $\pm e$ *(in the case* $p - q = 1 \mod 4$*) or* $\pm ie$ *(in the case* $p - q = 3 \mod 4$*) or* $\pm e^{1\dots n}$ *(in both cases).*

Then for any point $x_0 \in \Omega$ *there exists* $\varepsilon > 0$ *and there exists a function* $T = T(x) : O_\varepsilon(x_0) \to \mathcal{C}\ell^{\mathbb{C}}(p,q)$ *such that*

1. $T(x)$ *– a function of class* $C^k(O_\varepsilon(x_0))$*;*
2. $T(x)$ *– an invertible element of Clifford algebra* $\mathcal{C}\ell^{\mathbb{C}}(p,q)$ *for any* $x \in O_\varepsilon(x_0)$*;*
3. (a) $e^a = T^{-1}(x)h^a(x)T(x) \quad \Leftrightarrow \quad h^{1\dots n} = e^{1\dots n}$,
   (b) $e^a = -T^{-1}(x)h^a(x)T(x) \quad \Leftrightarrow \quad h^{1\dots n} = -e^{1\dots n}$,
   (c) $e^a = e^{1\dots n}T^{-1}(x)h^a(x)T(x) \quad \Leftrightarrow \quad h^{1\dots n} = e$,
   (d) $e^a = -e^{1\dots n}T^{-1}(x)h^a(x)T(x) \quad \Leftrightarrow \quad h^{1\dots n} = -e$,
   (e) $e^a = ie^{1\dots n}T^{-1}(x)h^a(x)T(x) \quad \Leftrightarrow \quad h^{1\dots n} = ie$,
   (f) $e^a = -ie^{1\dots n}T^{-1}(x)h^a(x)T(x) \quad \Leftrightarrow \quad h^{1\dots n} = -ie$,

   where equalities hold for $a = 1,\dots,n$ *and* $\forall x \in O_\varepsilon(x_0)$*;*

4. *The function $T(x)$ is defined up to multiplication by elements $\lambda(x)e+\nu(x)e^{1\ldots n}$, where $\lambda(x)$ and $\nu(x)$ - are complex functions of class $C^k(O_\varepsilon(x_0))$ such that element $\lambda(x)e+\nu(x)e^{1\ldots n}$ is invertible for any point of domain $O_\varepsilon(x_0)$.*

Proof. Proofs of Theorems 2 and 3 are similar to the proof of Theorem 1. We must use generalized Pauli's theorems [1] for Clifford algebras of odd dimension n. We consider expressions (instead of the expressions (2))

$$\sum_{A\in\mathcal{I}_{\mathrm{Even}}} h^A(x)F\gamma_A,$$

where sum is over multi-indices of even length $\mathcal{I}_{\mathrm{Even}}=\{A,\,|A|-\mathrm{even}\}$.

Element F is always among the basis elements $\{e^B\}$ or among the expressions $\{e^B+e^C\}$ of sum of two basis elemebts. So, F does not depend on the point x.

All other considerations are similar to considerations for the case of even n. The theorems are proved.

Theorems 2 and 3 evidently can be generalize to the case when both sets of Clifford algebra elements depend on points x of the Euclidian space.

References

[1] Shirokov D.S., Extension of Pauli's theorem to Clifford algebras, Dokl. Math., 84, 2, 699-701 (2011).

[2] Marchuk N.G., Shirokov D.S., Unitary spaces on Clifford algebras, Advances in Applied Clifford Algebras, Volume 18, Number 2, pp.237-254 (2008).